

Brazil

LEGEND

- Previously sampled Malaise trap site

Ecoregion

- Alto Paranã Atlantic forests
- Amazon–Orinoco–Southern Caribbean mangroves
- Araucaria moist forests
- Atlantic Coast restingas
- Bahia coastal forests
- Bahia interior forests
- Beni savanna
- Brazilian Atlantic dry forests
- Caatinga
- Caatinga Enclaves moist forests
- Campos Rupestres montane savanna
- Caqueta moist forests
- Cerrado
- Chiquitano dry forests
- Dry Chaco
- Fernando de Noronha–Atol das Rocas moist forests
- Guianan Highlands moist forests
- Guianan lowland moist forests
- Guianan piedmont moist forests
- Guianan savanna
- Gurupa várzea
- Humid Chaco
- Iquitos várzea
- Japurá–Solimões–Negro moist forests
- Juruã–Purus moist forests
- Madeira–Tapajós moist forests
- Marajó várzea
- Maranhão Babaçu forests
- Mato Grosso tropical dry forests
- Monte Alegre várzea
- Negro–Branco moist forests
- Northeast Brazil restingas
- Pantanal
- Pantepui forests & shrublands
- Pernambuco coastal forests
- Pernambuco interior forests
- Purus–Madeira moist forests
- Purus várzea
- Rio Negro campinarana
- Serra do Mar coastal forests
- Solimões–Japurá moist forests
- Southern Atlantic Brazilian mangroves
- Southern Cone Mesopotamian savanna Southwest
- Amazon moist forests
- Tapajós–Xingu moist forests
- Tocantins/Pindare moist forests
- Trindade–Martin Vaz Islands tropical forests
- Uatumã–Trombetas moist forests
- Uruguayan savanna
- Xingu–Tocantins–Araguaia moist forests